



**UNIVERSITI PUTRA MALAYSIA**

**EPIDEMIOLOGY AND DIAGNOSIS OF HUMAN LEPTOSPIROSIS IN  
MALAYSIA**

**ISAM MOHAMED ALI MOHAMED EL-JALII**

**IB 2000 2**

**EPIDEMIOLOGY AND DIAGNOSIS OF HUMAN LEPTOSPIROSIS IN  
MALAYSIA**

**By**

**ISAM MOHAMED ALI MOHAMED EL-JALII**

**Thesis Submitted in Fulfilment of the Requirements for the Degree of  
Doctor of Philosophy in the Institute of Bioscience  
Universiti Putra Malaysia**

**September 2000**



## **DEDICATION**

**TO THE MEMORY OF MY MOTHER, FATHER AND NEPHEW  
KHATAB**

**TO MY WIFE EMTENAN AND DAUGHTER RAYAN**

**TO MY BROTHERS AND SISTERS**

**TO ALL OF THEM WITH LOVE AND GRATITUDE**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia  
in fulfilment of the requirements for the degree of Doctor of Philosophy

**EPIDEMIOLOGY AND DIAGNOSIS OF HUMAN  
LEPTOSPIROSIS IN MALAYSIA**

By

**ISAM MOHAMED ALI MOHAMED EL-JALII**

**September 2000**

**Chairman: Prof. Dr. Abdul Rani Bahaman**

**Institute of Bioscience**

Retrospective study of human leptospirosis in Malaysia based on microscopic agglutination test (MAT) showed 13% overall prevalence of infection for the period 1983-1998. Results indicated that the prevalence was decreasing in the last five years (1994-1998). The prevalence was highest among Indians (16.67%) followed by Malays (11.48%) and the least among Chinese (5.88%). The 20-29 year-old group showed the highest prevalence of infection (17.13%). Less than 10 year-old group showed the least prevalence of infection (5.66%). Generally, many of the cases occurred between the ages of 20-50 years. Serological survey based on enzyme-linked immunosorbent assay (ELISA) showed a high overall prevalence (12.56%) of leptospiral infection. Kuala Lumpur showed the highest prevalence (19%) whilst Penang recorded the lowest prevalence (6.67%). No significant differences in the prevalence between the other states was noted.

A comparative study of three serological tests, namely enzyme-linked immunosorbent assay (ELISA), microscopic agglutination test (MAT) and indirect hemagglutination (IHA), test was carried out to evaluate these tests in the diagnosis of human leptospirosis. A total of 3000 serum samples from three groups of people were examined. In Group I, IgM and IgG-ELISA were able to detect a number of cases in the first sampling before MAT titres were detectable. In the second sampling, all samples positive for MAT were also positive for IgM-ELISA. IHA test gave positive reactions with only 38% of the samples while all samples were positive for ELISA. In Group II, ELISA detected IgM and IgG to leptospires in the samples which were negative to MAT. These were samples from patients with clinical signs of leptospirosis.

The polymerase chain reaction (PCR) was evaluated as a tool for diagnosis of leptospirosis and differentiation of leptospiral strains. Urine samples with as little as 10 serovar *hardjo* cells per ml of urine were positive on PCR indicating high sensitivity of the test. Detection of small number of leptospiral cells in urine by PCR was an advantage over culture. Random amplification polymorphic DNA (RAPD) fingerprinting was applied to differentiate leptospiral strains. Two primers were tested for their abilities to generate individual RAPD fingerprints. The DNA profiles obtained with each primer were distinct and reproducible. The fingerprint obtained could be useful for distinguishing the serovars up to the strain level. Profiles obtained revealed genetic heterogeneity between serovars belong to one serogroup.

Analysis of leptospiral DNA with restriction enzymes, *Hind* III, *Bam*H I and *Eco*R I revealed a high heterogeneity between the serovars examined. This high heterogeneity may be due to the large genome of the genus *Leptospira*. No relationship was found between the restriction patterns and the species from which the isolate was isolated. Similarities were observed among isolates of the same serovar. The 10 leptospiral field isolates were assayed for presence of plasmid DNA. Only two isolates were found to harbour plasmid DNA. The plasmid profiling obtained is of limited epidemiological value for differentiation of leptospiral isolates.

It appears that human leptospirosis is an endemic infection in Malaysia. The findings showed that ELISA was a suitable serological test for diagnosis of leptospirosis compared to MAT and IHA tests. On the other hand, the application of PCR and REA in the diagnosis of leptospirosis would be useful.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

## **EPIDEMIOLOGI DAN DIAGNOSIS LEPTOSPIROSIS DI MALAYSIA**

**Oleh**

**ISAM MOHAMED ALI MOHAMED EL-JALII**

**September 2000**

**Pengerusi: Prof. Dr. Abdul Rani Bahaman**

**Institut Biosains**

Kajian retrospektif terhadap leptospirosis manusia di Malaysia berasaskan ujian pengaglutinatan mikroskopi (MAT) menunjukkan prevalens jangkitan keseluruhan 13% untuk tempoh 1983 - 1998. Hasil kajian menunjukkan prevalens ini semakin kurang dalam tempoh lima tahun terakhir (1994-1998). Prevalens ini paling tinggi di kalangan kaum India (16.67%) diikuti kaum Melayu (11.48%) dan paling sedikit di kalangan kaum Cina (5.88%). Kumpulan umur 20-29 menunjukkan prevalens jangkitan paling tinggi (17.13%). Kumpulan umur kurang daripada 10 tahun menunjuk prevalens paling rendah (5.66%). Umumnya, banyak daripada kes ini berlaku pada kumpulan umur 20 - 25 tahun. Tinjauan serologi berasaskan asai imunoserap terangkai enzim (ELISA) menunjukkan prevalens jangkitan leptospira keseluruhan tinggi (12.56%). Kuala Lumpur menunjukkan prevalens paling tinggi (19%), manakala Pulau Pinang merekodkan prevalens paling rendah (6.67%). Tiada kelainan tererti wujud di antara negeri lain.

Suatu kajian perbandingan terhadap tiga ujian serologi, iaitu asai imunoerap terangkai enzim (ELISA), ujian pengaglutinatan mikroskopi (MAT), dan ujian penghemaglutinatan tak langsung (IHA) telah dijalankan untuk menilai ujian ini dalam diagnosis leptospirosis manusia. Sejumlah 3000 sampel serum manusia telah diperiksa. Dalam kumpulan I, IgM- dan IgG-ELISA dapat mengesan beberapa kes dalam pensampelan pertama, sebelum titer MAT dapat dikesan. Dalam pensampelan kedua, kesemua sampel yang positif dengan MAT juga positif dengan IgM-ELISA. Ujian IHA memberikan tindak balas positif untuk 38% daripada sampel sahaja, sambil kesemua sampel positif dengan ELISA. Dalam kumpulan II, ELISA mengesan IgM dan IgG terhadap leptospira dalam sampel yang negatif dengan MAT. Ini ialah daripada pesakit yang menunjukkan petanda klinikal untuk leptospira.

Tindak balas berangkai polimerase (PCR) telah dinilai sebagai alat untuk diagnosis leptospirosis dan untuk pembezaan strain leptospira. Sampel urin yang mengandungi hanya 10 *hardjo* sel per ml urin masih positif dalam ujian PCR, menunjukkan tingginya kepekaan ujian tersebut. Pengesanan sejumlah sel yang begitu kecil dalam urin melalui PCR memberi ujian ini kelebihan berbanding pengkulturan. Penyidikjarian DNA polimorfus penguatan rawak (RAPD) telah diguna untuk membezakan strain leptospira. Dua primer telah diuji untuk keupayaannya menjanakan sidikjari RAPD individu. Profil DNA yang diperolehi daripada setiap primer ini adalah nyata dan boleh dihasil semula. Sidikjari yang diperolehi mungkin



berguna untuk membeza serovar sehingga pada aras strain. Profil diperolehi menunjukkan keheterogenan genetik di antara serovar daripada satu serokumpulan.

Analisis DNA leptospira dengan enzim pengehadan, *Hind* III, *Bam*H I dan *Eco*R I menunjukkan keheterogenan tinggi di antara serovar yang diperiksa. Heterogenan tinggi ini mungkin disebabkan oleh genom besar pada genus *Leptospira*. Tiada perkaitan yang ditemui di antara pola pengehadan dan spesies daripada mana pencilan itu diperolehi. Persamaan telah dicerap di kalangan pencilan yang serovarnya sama. Pencilan 10 leptospira liar telah diasakan untuk kewujudan DNA plasmid. Hanya dua pencilan telah ditemui mengandungi DNA plasmid. Pemprofilan plasmid yang diperolehi itu mempunyai nilai epidemiologi terhadap dalam pembezaan pencilan leptospira.

Nampaknya leptospirosis manusia itu adalah suatu jangkitan endemik di Malaysia. Penemuan ini menunjukkan ELISA itu merupakan ujian serologi yang sesuai untuk diagnosis leptospirosis berbanding ujian MAT dan IHA. Sebaliknya, penggunaan PCR dan REA dalam diagnosis leptospirosis adalah memuaskan.

## ACKNOWLEDGEMENTS

First, my praise to Almighty Allah for giving me the power and will to complete this study.

My deepest gratitude and appreciation to my supervisor, Professor Dr. Abdul Rani Bahaman, Chairman of my Supervisory Committee, for his kind help, guidance, encouragement and invaluable assistance during the course of this study.

I would like to thank and express my deepest appreciation to Dr. Mohd. Azmi Mohd. Lila and Dr. Abdul Rahim Mutalib, members of my Supervisory Committee, for their invaluable advice and comments.

I wish to express my sincere gratitude to the Director of the Institute of Bioscience, former and present Deans of the Faculty of Veterinary Medicine, Universiti Putra Malaysia for allowing me to use the facilities of their laboratories during the course of this study.

I would like to thank Dr. Mangalam Sinniah from Institute for Medical Research (IMR), Kuala Lumpur for providing some of the serum samples. I am also grateful to Dr. Nadzri Salim from Faculty of Veterinary Medicine, Universiti Putra Malaysia for his help and advice.

I am most grateful to L. D. Smythe from WHO/FAO Collaboration Centre of Reference and Research on Leptospirosis, Queensland, Australia and Dr. C. A. Bolin from U. S. Department of Agriculture, USA for providing me with field and reference leptospiral strains used in this study.

I would also like to thank the staff members of Bacteriology Laboratory, Mr. Hajariah Salamah and Mr. Hamzah Adam who gave invaluable technical assistance. At the same time, I am very grateful to Mr. Zaid Othman, M. Munisamy and Omar Mohaideen for providing transportation during sample collection.

I would never forget those good friends and colleagues from whom I received kind assistance and help. I would like to thank Dr. Khairani Bejo, Dr. Abdul Wahid Saeed, Dr. Muthafar Elhaddawi, Dr. Izzeldin Babiker, Dr. Omer Arabi, Puan Zunita Zakaria, Dr. Awad El-Karim Mohd Bakhiet, Waleed Awad, Tongted Phumoonna, Dr. Md. Firoz Main. Dr. Karim EL-Jashamy and those who contributed in the completion of this study.

I am most grateful to the University of Khartoum (Sudan) for financial support. The research was wholly supported by the IRPA Grant (No. 06-02-04-055) obtained from the Ministry of Science, Technology and Environment (Malaysia).



Finally, I wish to express my deepest appreciation to my wife Emtenan, daughter Rayan and brothers Eltaib Saeed and Galal El-Jalii for their moral support and encouragement.

I certify that an Examination Committee met on 22nd September 2000 to conduct the final examination of Isam Mohamed Ali Mohamed El-Jalii on his Doctor of Philosophy thesis entitled "Epidemiology and Diagnosis of Human Leptospirosis in Malaysia" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Mohamed Shariff Mohamed Din  
Professor,  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Chairman)

Abdul Rani Bahaman, Ph.D  
Professor,  
Institute of Bioscience  
Universiti Putra Malaysia  
(Member)

Mohd. Azmi Mohd. Lila, Ph.D,  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

Abdul Rahim Mutalib, Ph.D,  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

Ansary Ahmad, Ph.D,  
Professor,  
Universiti Malaya  
(Independent Examiner)

  
\_\_\_\_\_  
MOHD. GHAZALI MOHAYIDIN, Ph.D,  
Professor/ Deputy Dean of Graduate School,  
Universiti Putra Malaysia

Date: 24 OCT 2000

This thesis submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Doctor of Philosophy.



---

KAMIS AWANG, Ph.D,  
Associate Professor,  
Dean of Graduate School,  
Universiti Putra Malaysia

Date: **14 DEC 2000**

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or currently submitted for any other degree at UPM or other institutions.



---

ISAM MOHAMED ALI MOHAMED EL-JALII

Date: 24. 10. 2000

## TABLE OF CONTENTS

	Page
DEDICATION.....	2
ABSTRACT.....	3
ABSTRAK.....	6
ACKNOWLEDGEMENTS.....	9
APPROVAL SHEETS.....	12
DECLARATION FORM.....	18
LIST OF TABLES.....	18
LIST OF FIGURES.....	20
LIST OF ABBREVIATIONS.....	22
 <b>CHAPTER</b>	
<b>I INTRODUCTION.....</b>	<b>24</b>
 <b>II LITERATURE REVIEW.....</b>	<b>29</b>
Leptospire.....	29
Epidemiology of Human Leptospirosis.....	33
Transmission of Leptospirosis.....	35
Pathogenesis of Leptospirosis.....	35
Immunity to Leptospirosis.....	37
Clinical Signs and Symptoms of Leptospirosis.....	39
Laboratory Diagnosis of Leptospirosis.....	41
Microscopic Examination.....	42
Histological Staining Techniques.....	43
Culture.....	44
Microscopic Agglutination Test (MAT).....	45
Enzyme-Linked Immunosorbent Assay (ELISA).....	46
Indirect Haemagglutination (IHA) Test .....	49
Complement Fixation Test (CFT).....	51
DNA Based Methods.....	51
Leptospiral Plasmids.....	63
Human Leptospirosis in Malaysia.....	64
Animal Leptospirosis in Malaysia.....	72
 <b>III EPIDEMIOLOGY OF HUMAN LEPTOSPIROSIS IN MALAYSIA.....</b>	<b>73</b>
Introduction.....	73
Materials and Methods.....	76
Retrospective Study.....	76
Serological Survey.....	77
Statistical Analysis.....	83
Results.....	83



	Retrospective Study.....	83
	Serological Survey.....	92
	Discussion.....	95
<b>IV</b>	<b>COMPARISON BETWEEN ENZYME-LINKED IMMUNOSORBENT ASSAY (ELISA) , MICROSCOPIC AGGLUTINATION TEST (MAT) AND INDIRECT HAEMAGGLUTINATION (IHA) TEST FOR DIAGNOSIS OF HUMAN LEPTOSPIROSIS.....</b>	<b>104</b>
	Introduction.....	104
	Materials and Methods.....	108
	Human Test Sera.....	108
	Microscopic Agglutination Test (MAT).....	110
	Enzyme-linked Immunosorbent Assay (ELISA).....	111
	Indirect Haemagglutination (IHA) Test.....	112
	Cross Reactivity.....	113
	Statistical Analysis.....	114
	Results.....	114
	Group I Samples.....	114
	Group II Samples.....	120
	Group III Samples.....	122
	Overall IgM and IgG-ELISA for the Different Groups.....	123
	ELISA IgM Titres to Leptospire for Group I samples.....	124
	ELISA IgG Titres to Leptospire for Group I Samples.....	125
	Discussion.....	127
<b>V</b>	<b>POLYMERASE CHAIN REACTION FOR DETECTION OF LEPTOSPIRES IN URINE AND DIFFERENTIATION OF LEPTOSPIRAL SEROVARS.....</b>	<b>136</b>
	Introduction.....	136
	Materials and Methods.....	140
	Leptospiral Strains.....	140
	Preparation of Leptospiral DNA.....	141
	Determination of DNA Concentration and Purity.....	143
	Urine Samples.....	143
	Preparation of Urine Samples for Specific PCR.....	143
	Specific PCR Primers.....	144
	RAPD-PCR Primers.....	145
	Specific PCR.....	146
	RAPD-PCR.....	146
	Gel Electrophoresis of Amplification Products.....	147
	Results.....	147
	Specific PCR.....	147
	RAPD-PCR.....	148
	Discussion.....	160

<b>VI</b>	<b>APPLICATION OF RESTRICTION ENDONUCLEASES ANALYSIS IN THE IDENTIFICATION OF LEPTOSPIRAL ISOLATES .....</b>	<b>165</b>
	Introduction.....	165
	Materials and Methods.....	167
	Leptospiral Isolates.....	167
	Preparation of Chromosomal Leptospiral DNA.....	168
	Determination of DNA Concentration and Purity.....	168
	DNA Digestion with Restriction Endonucleases.....	169
	Agarose Gel Electrophoresis and Photography of Digested Products.....	169
	Molecular Size Estimation of DNA Fragments.....	170
	Purification of Plasmid DNA.....	170
	Agarose Gel Electrophoreses of Plasmid DNA.....	171
	Results.....	172
	Digestion with Restriction Enzymes.....	172
	Plasmid DNA.....	179
	Discussion.....	184
<b>VII</b>	<b>GENERAL DISCUSSION AND CONCLUSION.....</b>	<b>188</b>
	<b>BIBLIOGRAPHY.....</b>	<b>200</b>
	<b>APPENDICES.....</b>	<b>223</b>
	Appendix A: Buffers Used for ELISA.....	224
	Appendix B: Preparation of Johnson and Seiter (JS) Medium...	227
	Appendix C: Reagents Supplied for IHA Kit.....	231
	Appendix D: Buffers and Reagents for Molecular Biology	233
	<b>BIODATA Of AUTHOR.....</b>	<b>237</b>

## LIST OF TABLES

Table	Page
2.1 Leptospiral serovars that have been isolated from man and animals in Malaysia.....	31
3.1 Prevalence of leptospirosis according to different states (1990-1998).....	85
3.2 Distribution of positive cases according to races.....	87
3.3 Prevalence of leptospirosis amongst different races.....	88
3.4 Distribution of positive cases of leptospiral infection (1988-1998) according to age groups.....	89
3.5 Clinical manifestations of positive cases of leptospiral infection (1988-1998).....	91
3.6 Distribution of IgM, IgG and overall ELISA positive samples amongst different states.....	93
3.7 IgM and IgG titres to leptospires for ELISA positive samples.....	94
4.1 Distribution of sera amongst different groups.....	109
4.2 Infecting serovars amongst MAT positive patients of Group I.....	115
4.3 Distribution of MAT titres during first and second samplings.....	116
4.4 IgM-ELISA during first and second samplings for Group I samples.....	117
4.5 IgG-ELISA during first and second samplings for Group I samples.....	118
4.6 Comparison between IgM, IgG-ELISA and IHA Test.....	119
4.7 IgM-ELISA during first and second samplings for Group II samples.....	121
4.8 IgG-ELISA during first and second samplings for Group II samples.....	121

4.9	IgM and IgG-ELISA for Group III samples.....	122
4.10	Overall IgM and IgG-ELISA for Group 1, 11 and 111 samples....	123
4.11	IgM titres to leptospire for Group I samples during first and second samplings.....	124
4.12	IgG titres to leptospire for Group I samples during first and second samplings.....	126
5.1	Leptospiral reference strains used in the study.....	140
5.2	Leptospiral field isolates used in the study.....	141
5.3	Sequences of 20 oligonucleotides screened for RAPD-PCR analysis.....	145
5.4	Distribution of fingerprinting profiles of reference strains obtained by RAPD-PCR using primers OPA 3 and OPA 20.....	159
5.5	Distribution of fingerprinting profiles of field isolates obtained by RAPD-PCR using primers OPA 3 and OPA 20.....	159
6.1	Leptospiral field isolates and reference strains used in the study	168
6.2	Molecular size of leptospiral DNA samples estimated from the fragments generated by digestion with Hind III restriction endonuclease enzyme.....	174
6.3	Molecular size of leptospiral DNA samples estimated from the fragments generated by digestion with BamH I restriction endonuclease enzyme.....	176
6.4	Molecular size of leptospiral DNA samples estimated from the fragments generated by digestion with EcoR I restriction endonuclease enzyme.....	178

## LIST OF FIGURES

Figure	Page
3.1 Epidemiology of leptospirosis.....	75
3.2 Prevalence of human leptospirosis in Malaysia (1983- 1998).....	84
3.3 Prevalence of leptospirosis according to different states (1990-1998).....	85
3.4 IgM and IgG titres to leptospires for ELISA positive samples.....	94
4.1 Distribution of MAT titres during first and second samplings.....	116
4.2 IgM titres to leptospires for Group I samples during first and second samplings.....	125
4.3 IgG titres to leptospires for Group I samples during first and second samplings.....	126
5.1 Gel electrophoresis of PCR products obtained from pure genomic DNA of leptospiral field isolates.....	151
5.2 Gel electrophoresis of PCR products obtained from urine samples seeded with 10, 20, 50, 100, 200, 400 <i>hardjo</i> cells per ml urine.....	152
5.3 RAPD profiles of leptospiral reference strains obtained with primer OPA 3 and electrophoresed on 2% agarose gel.....	153
5.4 RAPD profiles of leptospiral field isolates obtained with primer OPA 3 and electrophoresed on 2% agarose gel.....	154
5.5 RAPD profiles of leptospiral field isolates obtained with primer OPA 3 and electrophoresed on 2% agarose gel.....	155
5.6 RAPD profiles of leptospiral reference strains obtained with primer OPA 20 and electrophoresed on 2% agarose gel.....	156
5.7 RAPD profiles of leptospiral reference strains obtained with primer OPA 20 and electrophoresed on 2% agarose gel.....	157
5.8 RAPD profiles of leptospiral field isolates obtained with primer OPA 20 and electrophoresed on 2% agarose gel.....	158

6.1	REA profiles of leptospiral field isolates after digestion with Hind III and electrophoresed on 0.7% agarose gel.....	180
6.2	REA profiles of leptospiral field isolates after digestion with BamH I and electrophoresed on 0.7% agarose gel.....	181
6.3	REA profiles of leptospiral field isolates after digestion with EcoR I and electrophoresed on 0.7% agarose gel.....	182
6.4	Agarose gel (0.7%) electrophoresis of plasmid DNA from leptospiral field isolates.....	183

## LIST OF ABBREVIATIONS

Ab	Antibody
ABTS	2,2-azino-bis (3-ethylbenzthiazoline-6-Sulfonic acid)
AP-PCR	Arbitrary primed polymerase chain reaction
Bp	Base pair
BSA	Bovine serum albumin
°C	Degree Celcius
CAAT	Cross agglutination absorption test
CFT	Complement fixation test
CHEF	Contour-clamped homogenous electric field
CSF	Cerebrospinal fluid
D.D	De-ionized distilled
dNTP	Deoxy-nucleotide triphosphate
DFM	Dark-field microscopy
DNA	Deoxyribonucleic acid
Dr.	Doctor
EDTA	Ethylene diamine tetra-acetone
ELISA	Enzyme-linked Immunosorbent assay
FAO	Food and Agriculture Organization
FAT	Fluorescent antibody test
g	Gram
g/l	Gram/litre
G+C	Guanine+Cytisine
H <sub>2</sub> O <sub>2</sub>	Hydrogen peroxide
IgA	Immunoglobulin A
IgG	Immunoglobulin G
IgM	Immunoglobulin M
IHA	Indirect haemagglutination
IMR	Institute for Medical Research
JS	Johnson and Seiter
Kbp	Kilobase pairs
K L	Kuala Lumpur
L	Leptospira
LPS	Lipopolysaccharide
M	Molar
MAT	Microscopic agglutination test
Mda	Megadalton
mM	Milimole
M.S	Molecular size
M.W	Molecular weight
N.S	Negri Sembilan
O.D	Optical density
PBS	Phosphate buffer saline
PCR	Polymerase chain reaction

pH	Hydrogen-ion concentration
PLGE	Pulse field gel electrophoresis
PUO	Pyrexia of unknown origin
RAPD	Random amplified polymorphic DNA
REA	Restriction endonuclease analysis
RNA	Ribonucleic acid
rpm	Round per minute
SAT	Slide agglutination test
SDS	Sodium dodecyl sulphate
U	Unit
UK	United Kingdom
UM	Universiti Malaya
UPM	Universiti Putra Malaysia
US	United States
USA	United States of America
UV	Ultraviolet
µg	Microgram
µl	Microlitre
µM	Micro mole
V	Volt
VRI	Veterinary Research Institute
V/V	Volume/volume
WHO	World Health Organization
%	Percent



## CHAPTER I

### INTRODUCTION

Leptospirosis, also known as “march fever” and “mud fever,” is an important zoonotic disease, with important veterinary and public health impact (Dikken and Kmety, 1978; Gussenhoven *et al.*, 1997; Marcos *et al.*, 1997). Leptospirosis is caused by *Leptospira interrogans*. Based on immunological tests, more than 200 leptospiral serovars have been identified. The serovars could be placed into 23 serogroups (Soltys, 1979; Woodward *et al.*, 1997; Chu *et al.*, 1998). In Malaysia, 38 serovars from 13 serogroups have been known to occur (Bahaman, 1988).

In livestock, the disease causes important economic losses. Although this disease is usually mild and often subclinical, it can lead to great losses due to abortions, stillbirths, infertility, mastitis, weak progeny, decreased milk production, and with certain leptospiral serovars, death (Songer *et al.*, 1983; Thiermann, 1984; Bey and Johnson, 1986).

Leptospiral serovars can infect mammals including man. However, the pathogenicity and clinical manifestations of the disease depend on the animal